

## CLINICAL AND LABORATORY CHARACTERISTICS OF HIV INFECTION WITH INTESTINAL PARASITOSIS (BLASTOCYSTOSIS)

*Achilova M.M.*

*Samarkand State Medical University*

Clinical and laboratory symptoms of blastocystosis in patients with HIV infection were studied in this article. The study was conducted on 30 patients treated in the Samarkand Regional Infectious Diseases Hospital in 2021-2023. General clinical, parasitological, biochemical, serological, molecular-genetic, instrumental methods were used in the research work. From the gastrointestinal tract in patients: nausea, loose stools or constipation, pain in the epigastric area, around the navel and in the left flank area were found more in the main group compared to the control group. One of the clinical manifestations of blastocystosis is allergic symptoms. In the studied patients, urticaria was observed in 1 (7.14%) patient, dermatosis in 2 (14.2%) patients, skin itching in 4 (28.5%) patients, and allergic rhinitis in 1 (7.14%) patient. The number of eosinophil indicators in 12 (85.7%) cases in laboratory tests was 5-8%. In addition, in order to evaluate the immunological status of patients, when the IgE level was determined by the IFT method, it was found that this indicator increased when HIV infection was mixed with blastocystosis.

**Key words:** HIV infection, intestinal parasitosis, blastocystosis, immunoglobulin E.

**Relevance of the topic.** HIV infection is a chronic infectious disease of a viral nature, which mainly damages the immune system, causes the development of opportunistic diseases and tumors, and ends with death if antiretroviral therapy is not carried out [Raimondo M. et al., 2017; Guaraldi G., 2019]. HIV infection remains one of the major health problems in the world today. This disease mainly affects people aged 14-50. At the same time, HIV infection is causing great damage to the entire world economy. In HIV infection, along with a decrease in the activity of the immune system, the central nervous system, respiratory and digestive tracts are often damaged [Pokrovsky V.V., 2013]. Of the intestinal parasites, Blastocystis is mainly found in cats. In addition, this parasite has been identified in insects and leeches. Blastocystis enters the human body through non-compliance with hygiene rules [Prodeus T.V., 2014]. Today, due to the increase in the number of people with immunodeficiency, the frequency of meeting Blastocystis spp, Cryptosporidium spp and Cyclospora spp along with pathogenic intestinal protozoa Lamblia intestinalis and Entamaeba hystolitica has also increased [Bartlett D., 2012; Taylor TH, 2016]. In recent years, among intestinal parasitoses, there has been an increased interest in studying the specific characteristics

of the parasite of *Blastocystis hominis* in the human body. Today, by applying molecular genetic methods to detect parasitosis, it was found that 9 out of 17 subtypes of blastocysts (1-9) are found in humans, 1-3 ST in humans and animals, and the rest mainly in animals [Bouguero N.V. i dr., 2011, 2012; Rule K.N., 2017]. According to the literature, *Blastocystis* spp is distributed among 1 billion people (10% developed countries, 80% developing countries) all over the world [Abdiev T.A., 2013, 2018]. According to data, blastocystosis in people with immune deficiency is often manifested by intestinal dysfunction and allergic manifestations, in rare cases it is found that it passes without clinical symptoms [Balint A., 2014; Tamalee R., 2014; Gavrilyuk T.V., 2015].

In the current literature, there is insufficient information on the specific characteristics of HIV infection with intestinal parasites. Currently, the high incidence of intestinal parasitosis among people is due to the wide distribution of this disease in nature, and insufficient laboratory detection capabilities remain a special problem. Studying the clinical and laboratory characteristics of HIV infection with intestinal parasitosis is important in optimizing the treatment of this pathology. The above shows the relevance of this study.

**The purpose of the study:** to study the clinical and laboratory specific features of blastocystosis and HIV infection.


**Research object and methods.** A study was conducted on 30 HIV-infected patients treated in the Samarkand regional clinical hospital for infectious diseases in 2021-2023. Patients were divided into 2 groups: group 1 (main group) included 14 patients with HIV infection and blastocystosis, and group 2 (control group) included 16 patients with HIV infection.

The main part of the total number of patients included in the study was detected in 28 (93.3%) patients in the 3rd stage of HIV infection, in 2 (6.67%) patients in the 4th stage of the disease (AIDS stage). 10 (33.3%) of the total patients were prescribed antiretroviral therapy (ARVT), 20 (66.7%) did not receive ARVT.

Patients were diagnosed with "HIV infection" based on IFT and immunoblot examination at the Samarkand Regional AIDS Center in accordance with Order No. 277 of 2018 of the Ministry of Health of the Republic of Uzbekistan. The number of CD4 cells was determined in the blood. The number of HIV RNA in blood plasma was determined using the "Votex Rothergy Monitor Test" system. Invasion of parasites was determined by coproscopy (native/yo-on drop method of Kato and Miur) and fecal formalin ether sedimentation method. Coprological examination was conducted at the L.M. Isaev branch of the Republican Center for Specialized Epidemiology, Microbiology, Infectious and Parasitic Diseases. In addition to parasitological examination, the polymerase chain reaction (PCR) method was used to identify blastocysts. The amount of immunoglobulin E in blood serum was determined by IFT.

**Research results:** In the mixed form of HIV infection in group 1 (meeting with blastocystosis), 12 (85.7%) patients had complaints related to damage to the gastrointestinal tract, such as nausea, loose stools or constipation, pain in the epigastric area, around the navel, and in the left flank area. Subfebrile temperature (fever) was observed in 10 patients of this group (71.4%). Patients in group 2 had 2 times less complaints from the gastrointestinal tract in the mono-infection form of the disease compared to group 1, and 2.2 times less cases of subfebrile fever (31.3%). Subfebrile fever occurred in patients with low immune status in both groups (SD4+ cells were 206-316 cells/ $\mu$ l in patients of group 1, and 337-446 cells/ $\mu$ l in patients of group 2).

According to the scientific literature, one of the clinical manifestations of blastocystosis is allergic symptoms. Group 1 patients had urticaria in 7.1%, dermatosis in 14.2%, skin itching in 28.5%, and allergic rhinitis in one patient. Dermatomycosis was detected in only one of the patients of group 2 (Fig. 1).



1 - picture. Meeting of clinical signs when HIV infection is followed by blastocystosis

In laboratory tests, the number of eosinophils in 85.7% of patients of group 1 was in the range of 5-8%. In the control group, eosinophils were detected in 56.2% of cases in the range of 4-6%. In addition, the amount of IgE was determined by the IFT method to evaluate the immunological status of patients. In comparison to the control group, it was found that this indicator increased in group 1 patients with mixed blastocystosis with HIV infection.

**Summary.** Patients with HIV infection with blastocystosis experience more gastrointestinal clinical symptoms. Allergic symptoms of blastocystosis are more

evident in the mixed form of the disease. An increase in the amount of eosinophils and IgE in the blood of patients is one of the relative indicators of intestinal parasitosis.

### Used literature

1. Абдиев Т.А., Саидахмедова Д.Б., Сувонкулов У.Т., Ахмедова М.Д., Саипов Ф.С., Махмудова Л.Б., Абдиев Ф.Т., Вахобов Т.А., Коваленко Д.А., Качугина Л.В., Анваров Ж.А. Паразитарные болезни человека в Узбекистане. – Ташкент, 2018.
2. Ачилова М. М., Байжанов А. К., Ярмухамедова Н. А. ОЦЕНКА БЕЗОПАСНОСТИ ВЫСОКОАКТИВНОЙ АНТИРЕТРОВИРУСНОЙ ТЕРАПИИ У БОЛЬНЫХ ВИЧ-ИНФЕКЦИЕЙ.
3. Бартлетт Дж, Талант Дж, Фам П. Клинические аспекты ВИЧ-инфекции. Мл Р. Валент, 2012. - 528 с.
4. Байжанов А.К., Хикматуллаева А.С., Ачилова М.М. Лямблиоз ва бластоцистоз аниқланган беморларда ОИВ инфекциясининг кечиши / Вестник Ташкентской медицинской академии. - Ташкент, 2023. - №3/1. - С. 62-65. (14.00.00; №13).
5. Байжанов А.К., Ачилова М.М., Эргашева М.Я., Вахобова Н.Ш. Ичак паразитозларини специфик даволашнинг ОИВ инфекцияси кечишига таъсири // Вестник Ташкентской медицинской академии Спецвыпуск посвящён международной научно-практической конференции «Современные научные исследования в медицине: актуальные вопросы, достижения и инновации». - Ташкент, 2022. - С. 53-56.
6. Бугеро Н.В., Немова И.С., Потатуркина-Нестерова Н.И. Факторы персистенции простейшей фекальной флоры при дисбиозе кишечника // Вестник новых медицинских технологий. - 2011. - Т. 18. - №3.
7. Бугеро Н.В., Потатуркина-Нестерова Н.И. Результаты определения вирулентности *Blastocystis* spp. Методом рестрикционного анализа ДНК простейших // Фундаментальные исследования. - 2012. - №11-15.
8. Кучкарова Н.М., Байжанов А.К., Хикматуллаева А.С., Ачилова М.М., Абдукадырова М.А. ОИВ инфекциясида турли патологик жараёнларда ичак протозоозларининг ўрни // Сборник тезисов международной научно-практической конференции «Современные тенденции развития инфектологии, медицинской паразитологии, эпидемиологии и микробиологии». - Ургенч, 2023. - С. 124.
9. Покровский В.В. ВИЧ-инфекция и СПИД (Национальное руководство) - 2013. - С. 608.

10. Продеус Т.В., Федянкина Л.В., Фролова А.А. Морфологическая идентификация бластоцист // Медицинская паразитология и паразитарные болезни. – 2014. – № 1. – С. 9-13.

11. Balint A. Do not forget the stool examination! - cutaneous and gastrointestinal manifestations of Blastocystis sp. infection // Parasitology research. - 2014. - Т. 113. - №4. - С. 1585-1590.

12. Bayjanov A. K. et al. OIV infeksiyali bemorlarda ichak parazitozlarining tarqalganligi va blastotsistalarni genotiplash natijalari // Ўзбекистон Республикаси Санитария-эпидемиология ва жамоат саломатлиги хизмати журнали. – 2023. – Т. 3. – №. 3-4.

13. Bayjanov A.K., Achilova M.M., Ibadova N.X. The frequency of comorbid and opportunistic pathologies in patients with HIV infection // International scientific and practical conference «Cutting-edge science». - Shawnee, 2023. - P. 8-10.

14. Bayjanov A.K., Achilova M.M., Ibadova N.X. The frequency of comorbid and opportunistic pathologies in patients with HIV infection // International scientific and practical conference «Cutting-edge science». - Shawnee, 2023. - P. 8-10.

15. Kadirovich B. A. et al. RELATIONSHIP OF INCIDENCE OF INTESTINAL PARASITOSIS WITH THE LEVEL OF IMMUNODEFICIENCY IN PATIENTS WITH HIV INFECTION // Asian journal of pharmaceutical and biological research. – 2023. – Т. 12. – №. 2.

16. Mirkhamzaevna A. M., Yakubovna E. M., Shakhobidinovna V. N. Safety Assessment of Highly Active Antiretroviral Therapy in Patients with HIV Infection // EUROPEAN JOURNAL OF INNOVATION IN NONFORMAL EDUCATION. – 2022. – Т. 2. – №. 1. – С. 289-292.

17. Raimondo M., Camoni L., Suligoi B. et al. HIV-positive individuals on ART and with viral load suppressed in 12 Infectious Diseases Clinics in Italy: successes and disparities in the HIV Continuum of Care / AIDS Res Hum Retroviruses. 2017. – P. 1056-61. Rule K.N. Blastocoel cell-free DNA, a marker of embryonic quality / Fertil. Steril. – 2017. – V. 108. – № 3. – P. e106.